



Impetigo

Impetigo is a skin infection caused by either Staphylococcus aureus (staph) bacteria or Group A Streptococcus (strep) bacteria. It is more common in children than adults, and most often occurs in the summer. It is usually caused when bacteria enter the body through breaks in the skin, such as scrapes, cold sores, insect bites or patches of eczema.

Symptoms

- + A child with impetigo will have a skin rash that looks like a group of small blisters or red bumps. After these blisters appear they often burst, and fluid seeps out. When the fluid dries, the blisters become coated with a yellow or grey crust. The blistered area will often be surrounded by redness. The rash will spread if it is not treated.
- + This rash usually appears around the nose, mouth, and other parts of the face. It can also appear on any skin not covered by clothes, such as arms and legs.
- In severe cases of impetigo there may be swelling of the lymph glands in the face or neck, severe pain, worsening redness, weakness and fever.

IS IMPETIGO SERIOUS?

Impetigo is usually a mild infection. Children with impetigo will have little discomfort other than occasional itching. In rare cases, the

bacteria that cause impetigo can get into the bloodstream and cause kidney disease.

HOW IS IT SPREAD?

Impetigo is very infectious and spreads easily from person-to-person through nasal discharge or skin-to-skin contact. For example, when someone touches the rash with their hand and then touches another person the infection can be passed on. Impetigo can also be spread by touching objects contaminated with the bacteria. It can take anywhere from 1 to 10 days for the rash to appear after you have been infected.

You should practice good personal hygiene when a family member or someone you know has impetigo. It is very important to wash your hands after touching the rash, as impetigo is easily spread by hand-to-hand contact.

A child with impetigo should not share towels, clothes, face cloths, bedding, toys, or personal items such as combs or brushes with anyone else.

WHEN TO GET HELP

See your family doctor if you think your child has impetigo.

Be careful to protect yourself and other children from close contact with children with impetigo. If your health care provider finds that your child does have impetigo, it can be easily

bc**emn**.ca

The information in this document is intended solely for the person to whom it was given by the health care team. This information does not replace the advice given to you by your health care provider.



Impetigo - Continued

treated with antibiotic cream. Oral antibiotics may be prescribed for more severe cases where the rash covers large areas of the body. It is important that your child keeps taking all of the prescribed antibiotics until they are used up, even if the rash has gone away.

With treatment, impetigo should be gone in about 7 days.

HOW CAN PARENTS HELP?

Keep your child at home.

Impetigo easily spreads at daycares, schools and summer camps. A child with impetigo should stay at home.

Limit contact with children with impetigo.

If you know of another child who has impetigo, you should watch for signs in your own children, and try to limit your child's contact with them.

Wash your hands.

If your child gets impetigo, everyone in your home will need to wash their hands carefully and often, especially after any physical contact.

Wash clothes and linens separately.

If your child has impetigo, it is important to wash their clothes and bed linens separately from other's. Use hot water to wash, and dry items in a hot dryer.

Tell your child not to touch or scratch sores.

Children with impetigo should be discouraged from touching or scratching their sores. Keep their fingernails short and clean. They should also be helped to wash their hands often.

As the disease develops, loose crusts on the skin can be soaked for 15 to 20 minutes with a warm wet facecloth, then gently washed off with soap and water and patted dry.

WHEN CAN MY CHILD RETURN TO SCHOOL OR DAYCARE?

Your child can usually return to school or daycare after 24 hours of antibiotic treatment. This will reduce the chance of spreading impetigo to other children.

RELEVANT RESOURCES

+ For more information on handwashing, see https://www.healthlinkbc.ca/healthlinkbc-files/hand-washing

TO LEARN MORE ASK

- + Your family doctor.
- HealthLinkBC call 8-1-1 (7-1-1 for deaf or hard of hearing) or go online to www.HealthLinkBC.ca

RECOMMENDED FOLLOW UP

Follow	up	with	your	primary	care	provider
in	day	S.				

Reproduced with permission from HealthLink BC.

Find this information sheet on the BC Emergency Medicine Network website:

www.bcemn.ca/clinical_resource/impetigo/

bc**emn**.ca

