

Section I	Scenario	Demographics
-----------	----------	--------------

Scenario Title:	Massive Upper GI Bleed	
Date of Development:	10/03/2015	
Target Learning Group:	□ Juniors (PGY 1 – 2) □ Seniors (PGY \ge 3)	🖂 All Groups

Section II: Scenario Developers

Scenario Developer(s):	Cheryl ffrench
Affiliations/Institution(s):	University of Manitoba
Contact E-mail (optional):	emerg13@gmail.com

Section III: Curriculum Integration

	Learning Goals & Objectives		
Educational Goal:	To manage a patient who presents to the ED with a massive UGIB.		
CRM Objectives:	1) Anticipate and appropriately plan for intubation in a patient with a significant		
	UGIB		
	2) Consider a massive transfusion strategy in a non-trauma context		
Medical Objectives:	1) Employ adjunctive medications in the treatment of an undifferentiated		
	massive UGIB including octreotide/somatostatin analogues, intravenous		
	proton pump inhibitor, and broad-spectrum antibiotics.		
	2) Prioritize airway management and Blakemore tube insertion for the		
	stabilization of a patient with a massive UGIB prior to definitive therapy.		
	3) Recognize the importance of limited crystalloid therapy and early blood		
	transfusion in the setting of hypovolemic shock secondary to UGIB.		

Case Summary: Brief Summary of Case Progression and Major Events

A 58-year-old male known for alcoholism presents to the emergency department with a two-day history of hematemesis with an active, massive upper GI bleed due to esophageal varices. The patient deteriorates into hypovolemic shock requiring medical management, blood transfusions, intubation for airway protection and insertion of a Blakemore tube before definitive management.

References

Marx, J. A., Hockberger, R. S., Walls, R. M., & Adams, J. (2013). *Rosen's emergency medicine: Concepts and clinical practice*. St. Louis: Mosby.
 Chavez-Tapia NC, Barrientos-Gutierrez T, Tellez-Avila FI, Soares-Weiser K, Uribe M. Antibiotic prophylaxis for cirrhotic patients with upper gastrointestinal bleeding. Cochrane Database of Systematic Reviews 2010, Issue 9.
 Chase, C. (2014, September 2). Management and Dispo of Upper GI Bleed - emdocs. Retrieved May 20, 2015, from http://www.emdocs.net/management-dispo-upper-gi-bleed/





Section VI: Scenario Progression

Scenario States, Modifiers and Triggers				
Patient State Patient Status Learner Actions, Modifiers & Triggers to Move to Next State				
1. Baseline State Rhythm: Sinus Tach HR: 115/min BP: 105/60	Alert, drowsy, intoxicated, GCS: 13 Active	Learner Actions - Monitors/Full vital signs - Physical exam	<u>Modifiers</u> Changes to patient condition based on learner action	
RR: 24/min O ₂ SAT: 96 % T: 37.8°C	intermittent Bright red blood hematemesis	- IV NS Fluid bolus - CXR/AXR - EKG - Blood work ordered, including type & screen + Coags	Triggers For progression to next state - 2 minutes → 2. Hemorrhagic Shock	
2. Hemorrhagic Shock HR → 140 over 30 seconds BP → 85/50	1 episode of copious hematemesis GCS: 7 (E1V3M4)	Learner Actions - IV NS Fluid bolus #2 - Blood Transfusion 2U - Massive Transfusion protocol activation - IV PPI (Bolus + infusion) - Central line access - Octreotide bolus + infusion - Intubation - +Vasopressin infusion	$\frac{\text{Modifiers}}{\text{-} Blood transfusion/MTP}$ initiation → HR 120, BP 95/60 - IV NS bolus w/o blood → HR 125, BP 90/55 transiently. $\frac{\text{Triggers}}{\text{-} Intubation} \rightarrow 3. Intubation$ - 5 minutes w/o intubation → 6. $PEA arrest$	
 3. Intubation <u>Display</u> (if possible) Vt: 500ml RR → 12 FiO2: 1.0 EtCO₂ = 50, normal waveform 4. Sengstaken-Blakemore tube 	Pt requires significant suction, able to intubate via direct laryngoscopy	Learner Actions - 2 suction tips prepared - Difficult Airway Kit - Appropriate induction med - Appropriate paralytic med - Sengstaken-Blakemore tube GI/ICU/Surgery consult Learner Actions - - Preparation (HOB at 45°, test balloons, check tube markings) - - Placement of tube - CXR - IV Antibiotics (Ceftriaxone)	Modifiers Triggers - Sengstaken-Blakemore tube - Sengstaken-Blakemore tube - 5 minutes without Blakemore tube placement after intubation → 6. PEA Arrest Modifiers Triggers - Blakemore tube confirmed with CXR → 5. Resolution.	
5. Resolution HR \rightarrow 110 BP \rightarrow 105/80 6. PEA Arrest	Pt transferred for emergent endoscopy.	- ± + PCC/Vit K - ± TXA	END SCENARIO END SCENARIO PRN	
HR \rightarrow 130 BP \rightarrow 0/0 O ₂ SAT \rightarrow 0		- CPR - Epinephrine	GI/ICU arrive and declare patient unsalvageable	



© 2015 EMSIMCASES.COM

This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.





General Debriefing Plan				
🗌 Individual	Group	🗌 With Video	🛛 Without Video	
	Objectives			
Educational Goal:	Educational Goal: To manage a patient who presents to the ED with a massive UGIB.			
CRM Objectives:	 Anticipate and appropriately plan for intubation in a patient with a significant UGIB Apply a massive transfusion strategy in a non-trauma context 			
Medical Objectives:	 1) Employ adjunctive medications in the treatment of a likely variceal UGIB including octreotide/somatostatin analogues and broad-spectrum antibiotics. 2) Prioritize airway management and Blakemore tube insertion for the stabilization of a patient with a massive UGIB prior to definitive therapy. 3) Recognize the importance of limited crystalloid therapy and early blood transfusion in the setting of hypovolemic shock secondary to UGIB 			
	Sample Questi	ons for Debriefing		
 What are the medical management options for a significant variceal bleeding? What are the evidence based outcomes for the use of octreotide in UGIB? Antibiotics? PPI? What is a massive transfusion protocol? When and how would you initiate an MTP for a non-trauma patient? What are the airway considerations in a patient with a massive UGIB? Describe the set up and insertion of a Sengstaken-Blakemore tube. Describe the coagulopathy associated with cirrhosis and its effect on management of the cirrhotic UGIB. 				
Key Moments				
- Hemorrhagic Shock				
- Intubation				
- Sengstaken-Blakemore tube insertion				

